| 1. | The | normal value of p | oH of | blood is: | | | | | |
|-----|--------|--------------------|---------|---------------|--------|---------|-------------------|---------|---------------|
| | (1) | 7.4 | (2) | 7.3 | | (3) | 7.5 | (4) | 7.2 |
| 2. | | mal bleeding time | | 2 4 | | (2) | 1 2 | (4) | 1 5 |
| | (1) | 1 - 2 minutes | (2) | 1 - 4 minu | tes | (3) | 1 - 3 minutes | (4) | 1 - 5 minutes |
| 3. | | he following are | , . | | - | | | | |
| | (1) | long bones | (2) | short bone | :S | (3) | flat bones | (4) | regular bones |
| 4. | Nor | mal urine output | for ar | adult duri | ng 24 | hours | is: | | |
| | (1) | 1500 ml | (2) | 1400 ml | | (3) | 1300 ml | (4) | 1200 ml |
| 5. | Initia | ation of milk secr | etion | is promoted | by: | | | | |
| | (1) | Leuteinizing ho | rmon | e | (2) | Folli | cle - stimulating | hormo | one |
| | (3) | Prolactin horm | one | | (4) | Som | atotrophic horm | one | |
| 6. | Food | d intoxication due | e to ca | nned food i | s caus | ed by | ·: | | |
| | (1) | staphylococcus | | | (2) | clost | ridium Botulinu | m | |
| | (3) | streptococcus | | | (4) | salm | onella | | |
| 7. | Glov | ves are sterilized | by aut | oclaving for | : | | | | |
| | (1) | 20 minutes at 2 | | | | | | | |
| | (2) | 30 minutes at 2 | | | | | | | |
| | (3) | 20 minutes at 2 | | | | | | | |
| | (4) | 30 minutes at 2 | 50° F | to 254° F | | | | | |
| 8. | | m inhalation in p | | with chest | - | | - | | |
| | (1) | provide comfor | | | (2) | | ove thick tenacio | ous spi | itum |
| | (3) | help in rest and | l sleep |) | (4) | relie | ve pain | | |
| 9. | | objectives of moi | | | | | • | | |
| | (1) | to soothen the | | | | | to relieve disco | | |
| | (3) | to stimulate the | infla | mmation pr | ocess | (4) | to reduce temp | oeratui | re |
| 10. | The | crutch walking g | gait th | at can bear t | the we | eight l | by the strong leg | ; is : | |
| | (1) | Tripod shufflin | | | (2) | | r point gait | | |
| | (3) | Three point gai | it | | (4) | Two | point gait | | |
| 11. | The | objective of mair | | | hanic | s whi | le providing nur | sing ca | are is to : |
| | (1) | make proper us | | | | | | | |
| | (2) | be physically a | | | _ | | | | |
| | (3) | maintain body | _ | | oid st | rain | | | |
| | (4) | maintain body | equili | brium | | | | | |

| 12. | Whi | | al trac | tion all the following should be kept in mind |
|--------|------------------|---------------------------------------|---------------|---|
| | (1) | remove the weight while changin | g the | position |
| | (2) | maintain the accuracy of weight | | _ |
| | (3) | ensure that weight does not touch | | |
| | (4) | remove the weight if required wi | thout | causing jerk |
| 13. | | 0.1 | e kep | t in mind while doing any sterile procedure |
| | $exce_{1}$ (1) | face the sterile field | | |
| | (2) | keep sterile equipment below the | waist | level |
| | (3) | never cross sterile area | | |
| | (4) | do not speak over sterile field | | |
| 14. | Whi | le feeling the Pulse when the beat i | s skip | ped it is referred as : |
| | (1) | lrregular pulse | (2) | Dicrotic pulse |
| | (3) | Intermittent pulse | (4) | Regular pulse |
| 15. | Whe | en the patient has discomfort in bre | eathing | g while standing is called as : |
| | (1) | cheyne - stokes respiration | (2) | Dyspnea |
| | (3) | stertorous | (4) | orthopnea |
| 16. | Lith | otomy position is given to a patien | t in al | l of the following conditions except: |
| | (1) | vaginal delivery | (2) | enema |
| | (3) | rectal examination | (4) | catheterization |
| | | | | |
| 17. | | verbal communication includes all | | • |
| | (1) | Facial expressions | (2) | Gestures Protection and Golden |
| | (3) | Vocabulary · | (4) | Posture and Gait |
| 18. | The | route used for Mantoux testing is: | : | |
| | (1) | Intramuscular | (2) | Intradermal |
| | (3) | Subcutaneous | (4) | Intravenous |
| 19. | Ahn | normal collection of fluid within a c | definit | e sac is known as |
| 1. / . | (1) | tumor (2) wart | acmin. | (3) cyst (4) polyp |
| | () | (=, | | (2) POSP |
| 20. | Mor | phine belongs to one of the follow | ing gr | oup of drugs : |
| | (1) | belladonna alkaloids | (2) | anticholinergics |
| | (3) | opiates | (4) | barbiturates |
| 21. | Leve | el of consciousness of a head injury | , patie | nt is assessed by a nurse using : |
| | (1) | head injury chart | (2) | Snellen's chart |
| | (3) | rapid assessment scale | (4) | glassgow coma scale |
| | , , | - | , , | |

| 22. | Patie | ent is given the following position | n durin | g abd | ominal surgery | : | |
|------------|-------|--|----------|---------------------------------------|-------------------|-----------|----------------------|
| | (1) | lithotomy | (2) | dors | al recumbent | | |
| | (3) | trendelendburg | (4) | Sim | s | | |
| | | | | | | | |
| 23. | Pleu | ral effusion is also known as : | | | | | |
| | (1) | pneumothorax | (2) | hem | othorax | | |
| | (3) | hydrothorax | (4) | spoi | ntaneous pneun | nothora | x |
| 24. | IPPB | stands for : | | | | | |
| | (1) | Immediate Positive Pressure Br | eathing | , | | | |
| | (2) | Intermittent Positive Pressure E | Breathin | ıg | | | |
| | (3) | Intermission Positive Pressure | Breathir | ng | | | |
| | (4) | Indirect Positive Pressure Breat | hing | | | | |
| 25. | All | of the following measures should | be tak | en du | ring blood trans | fusion : | reaction except : |
| | (1) | collect blood sample immediate | | | | | · |
| | (2) | stop transfusion | | | | | |
| | (3) | provide psychological support | | | | | |
| | (4) | monitor vital signs | | | | | |
| 26. | Patie | ent with sinus bradycardia will h | nave he | art ra | te below : | | |
| | (1) | 40 per minute | (2) | | er minute | | |
| | (3) | 60 per minute | (4) | | er minute | | |
| | ` / | | () | | | | • |
| 27. | | of the following approaches used is referred as: | to categ | gorize | the casualties du | ring di | saster for providing |
| | (1) | Triage (2) Triad | | (3) | Screening | (4) | Scrutinizing |
| 28. | The | immediate nursing intervention | for a na | tiont | with opictavic d | ue to b | aemodialveie ie : |
| 40. | | Allow the patient to breath no | _ | · · · · · · · · · · · · · · · · · · · | with epistaxis a | ac to It | acmodiary sis is: |
| | (2) | Give dorsal recumbent position | | | | | |
| | (3) | Call the Doctor | | | | | |
| | (4) | Pack the nose with paraffin ga | 1170 | | | | |
| 29. | ` ' | of the following are monitored by | | sa zuh | an a nationt is o | n diuro | tice avaget : |
| 29. | | | y a murs | | restoring fluid | | • |
| | (1) | assessing weight | -1. | (2) | | | |
| | (3) | maintaining intake output cha | 11 | (4) | monitoring vi | iai signs | 5 |
| 30. | | of the acute complications of dia | | | | | |
| | (1) | Hypoglycemia | (2) | | etic neuropathy | 7 | |
| | (3) | diabetic retinopathy | (4) | dial | etic foot | | |
| | | | | | | | |

| 31. | Com | plications of retro | pubic | prostatecto | omy is | ; : | | | | |
|-----|-------------|--|---------------|---------------------------------------|--------|----------------|--------------------|----------|-------------------|----|
| | (1) | recurrent obstru | ction | | (2) | diffi | cult to control h | emorrh | age | |
| | (3) | urinary leakage | | | (4) | incid | lence of increase | ed hem | orrhage | |
| 32. | | ires characterized of consciousness i | - | | d by a | abrup | t fall with jerky | moven | nents of body an | d |
| | (1) | partial seizure | | | (2) | psyc | chomotor seizure | 9 | | |
| | (3) | focal seizure | | | (4) | gene | eralized seizure | | | |
| 33. | | period between tl vn as : | he ent | ry of agent | of dis | s e ase | and before appe | arance | of clinical signs | is |
| | (1) | prodromal perio | ď | | (2) | prec | linical period | | | |
| | (3) | incubation perio | | | (4) | • | eptible period | | | |
| 34. | | l number of cases | perta | nining to an | y dise | ase e | xisting in a com | munity | at a given time | is |
| | (1) | prevalence rate | | | (2) | incid | dence rate | | | |
| | (3) | morbidity rate | | | (4) | mor | tality rate | | | |
| 35. | | ngle tablet (0.5g) o | | orine is suff | | | | | | |
| | ` / | 10 litres of wate | | | (2) | | tres of water | | | |
| | (3) | 20 litres of wate | ı. | | (4) | 25 li | tres of water | | | |
| 36. | Duri | ng early stages of | Xero | phthalmia t | the an | nount | of vitamin A ad | ministe | ered orally is: | |
| | (1) | 200,000 IU | (2) | 100,000 IU | | (3) | 50,000 IU | (4) | 300,000 IU | |
| 37. | The than | incidence of thyr | oid is | common v | when | the a | mount of iodine | in salt | consumed is le | ss |
| | (1) | 30 ppm | (2) | 15 ppm | | (3) | 20 ppm | (4) | 25 ppm | |
| 38. | Revi | sed National Hea | lth Po | olicy of Indi | a was | anno | ounced in : | | | |
| | (1) | 2000 | (2) | 2001 | | (3) | 2002 | (4) | 2003 | |
| 39. | New | born delivered in | n the i | hospital are | oiver | all o | of the following s | zaccines | e orcont · | |
| 0,1 | (1) | BCG | (2) | Hepatitis l | _ | (3) | Oral Polio | (4) | Hepatitis A | |
| 10 | Y | halian arai da 6 | 1.1.1 | | | | | | | |
| 40. | incu (1) | bation period of o 12 - 14 days | :nicke (2) | n pox 1s : 10 - 14 da [.] | vs | (3) | 14 - 16 days | (4) | 14 - 18 days | |
| | | • | , | | | | · | (-) | ,- | |
| 41. | | of the major com | - | | Mum | ps in | males is : | | | |
| | (1) | Weakness and f | atigue | ? | (2) | Ster | • | | | |
| | (3) | Ear-ache | | | (4) | Infe | rtility | | | |

| 42. | Trea | tment of pneumonia in infants und | ler 2 r | months of age is by giving : |
|-------------|-------|--|------------|-------------------------------------|
| | (1) | Injection Benzyl Penicillin and In | | 0,00 |
| | (2) | Injection Procain Penicillin and In | , | • |
| | (3) | Injection Benzyl Penicillin and In | , | |
| | (4) | Injection Procain Penicillin and I | - | • |
| 43. | DOT | S stands for : | | |
| | (1) | Directly Observed Treatment Sch | eme | |
| | (2) | Directly Observed Treatment Sho | rt Co | urse |
| | (3) | Directly Observed Therapeutic So | ervice | |
| | (4) | Directly Observed Tuberculosis Se | ervice | s |
| 44. | Actio | ons taken for the community prior | to occ | currence of disease is referred as: |
| | (1) | Rehabilitation | (2) | Specific protection |
| | (3) | Tertiary prevention | (4) | Primary Prevention |
| 45. | All c | of the following are principles of Pr | rimary | y Health Care <i>except</i> : |
| | (1) | advance technology | (2) | equitable distribution |
| | (3) | community participation | (4) | inter-sectora! coordination |
| 46. | Mate | ernal Mortality Rate of India in 201 | 1 is: | |
| | (1) | 258 per 100,000 live births | (2) | 254 per 100,000 live births |
| | (3) | 252 per 100,000 live births | (4) | 256 per 100,000 live births |
| 47 . | All o | of the following are community nu | tritior | n programmes for children except : |
| | (1) | Iron and folic acid supplementati | | |
| | (2) | Vitamin A prophylaxis | | |
| | (3) | Balwadi nutrition | | |
| | (4) | Midday meal | | |
| 48. | Whe | en a responsibility to a nursing offic | er is | given with authority it is called : |
| | | Delegation of power | | • |
| | (3) | Centralization of power | (4) | Span of control |
| 49. | The | objective of nursing supervision in | clude | all the following excent: |
| | (1) | to provide quality care to the pat | | in the remember of |
| | (2) | to improve performance of the m | | personnel |
| | (3) | to pinpoint the mistakes done | | |
| | (4) | to facilitate in work | | |
| 50. | The | best method for clinical teaching is | : | |
| | (1) | role play | (2) | nursing care plan |
| | (3) | symposium | (4) | seminar |
| | (-) | -/ . / | (3) | |

| 51. | Skills (1) (2) (3) (4) | s of a good nurse clearly defined to ignoring non-per explaining task a providing resour | he goarform | als to be ach ers and exp dvantages o | nieved ecting of outp | othe: | | | pt : | | | |
|-----|------------------------------------|--|---------------|---|-----------------------------|----------------------|------------------|-------------------------------------|----------|--------|----------------|---|
| 52. | Class (1) | sroom teaching is course plan | done (2) | | - | | (3) | lesson pla | n | (4) | unit plan | |
| 53. | All o (1) (3) | of the following ar to increase the k repeat all the pro | nowle | edge | | rescar (2) (4) | to pr | cept : ovide evide nd out new | | ased o | care | |
| 54. | While except | le conducting rese | earch | study on th | e pati | ents a | all of | the followir | ng sho | uld b | e considered | ł |
| | (1) (3) | informed conser freedom of harn | | | (2) (4) | | identi ire pa | ality rticipation | | | | |
| 55. | , , | se researcher shou | ıld rev | view all of t | | | • | , | ing a c | etudy | excent: | |
| 33. | (1) (3) | old research rep journals | | view all of t | (2) (4) | | publi | shed docum | 411 | study | ехсері . | |
| 56. | | se researcher can t cople : | ise on | e of the foll | owing | ; meth | ods fo | or collecting | ; data : | from l | arge numbe | r |
| | (1) | questionnaire | (2) | interview | | (3) | discı | ussion | (4) | telep | phonic talk | |
| 57. | The : (1) | sum of all the values median | ues di (2) | vided by th range | e total | l num (3) | ber is mod | | (4) | mea | n | |
| 58. | Beha | viours which get | s tran | smitted fror | n one | gene | ration | to another | is kno | own a | s : | |
| | (1) | customs | (2) | beliefs | | (3) | norn | ns | (4) | culti | ure | |
| 59. | The (1) | indices of standar Type of family | d of l | iving are al | l of th (2) | | | except : | | | | |
| | (3) | Goods one const | umes | | (4) | Serv | ices o | ne enjoys | | | | |
| 60. | | of the following i | s a de | efense mech | | | - | | l: | | | |
| | (1) (3) | Frustration Denial | | | (2) (4) | 45 45 | ression | n hdrawal | | | | |
| 61. | | n an individual is | not a | able to decid | đe amo | ongst | two e | qually imp | ortant | goals | it is referred | d |
| | as : (1) | Substitution | (2) | Conflict | | (3) | Repr | ression | (4) | Proj | ection | |

| 62. | Hier (1) | archy of needs w Maslow | as giv (2) | en by : Skinner | | (3) | Freud | (4) | Erickson |
|-----|---------------------------------|---|---------------------------|--|------------------|----------------------|--|-------------------------|-----------------------------|
| 63. | The | newborn should | alwav | s be kept cl | ose to | the n | nother for effec | tive : | |
| | (1) | attachment | (2) | parenting | | (3) | caring | (4) | bonding |
| 64. | All (1) | of the following a Id | re exp (2) | plained by F Grandiosi | | in Psy (3) | vchoanalytical Ego | theory <i>ex</i> (4) | <i>ccept</i> : Super ego |
| 65. | Indi (1) | an Nursing Coun 1946 | cil Ac (2) | t was passe 1947 | d in : | (3) | 1948 | (4) | 1949 |
| 66. | Whe (1) (3) | en nurses are assi functional meth team method | | specific task | s for (2) (4) | patie | f patients is re ent method nary care meth | | : |
| 67. | Dur (1) (2) (3) (4) | ing a performanc employee's curr career plan of t determining wh employees cont | rent le he em ny em | vel of perfo ployee ployee has p | rmano | ce med p | | cus on all | except : |
| 68. | List (1) (3) | of activities preso job specification job evaluation | | for various | (2) (4) | job (| professionals description summary | is knowr | n as: |
| 69. | One (1) | of the methods finduction | or sta | ff developm introducti | | sed in (3) | the new settin | ng is refer (4) | rred as : orientation |
| 70. | All (1) | of the following a democratic | are Lea (2) | adership sty idealistic | les <i>ex</i> | cept : (3) | autocratic | (4) | laissez faire |
| 71. | ABC (1) (3) | analysis for investoring criteria cost criteria | entory | control is b | (2) (4) | item | i criteria ing criteria | | |
| 72. | Adv (1) | rertisement and in recruitment | ntervio (2) | ewing for se planning | lectin | g new (3) | staff refers to marketing | (4) | staffing |
| 73. | | en ANS of one ur S of another unit vertical commu horizontal com | it is a inicati | n example o on | | diag | garding trainin gonal commun vard communi | ication | ing personnel with |

| 74. | Nun | nber of subordina | tes an | administra | tor ca | n con | trol or supervi | se is kno | wn as : |
|-----|--------------|---------------------------|--------|----------------|------------|--------|------------------------|-----------|--------------------|
| | (1) | unity of comma | nd | | (2) | span | of control | | |
| | (3) | delegation | | | (4) | hiera | archy | | |
| 75. | Infla | mmation of vagir | nal ep | ithelium ca | used l | oy can | dida albicans i | is known | as: |
| | (1) | monilial infection | n | | (2) | trich | omoniasis infe | ection | |
| | (3) | Bartholin's glan | d infe | ction | (4) | cerv | icitis | | |
| 76. | Infla | mmation of lining | g of u | terus is kno | wn as | 3; | | | |
| | (1) | vaginitis | | | (2) | endo | ocervicitis | | |
| | (3) | adenomyositis | | | (4) | endo | ometritis | | |
| 77. | All | of the following a | re the | high risk fa | actors | for b | reast cancer <i>ex</i> | cept : | |
| | (1) | first parity before | | - | (2) | | ly history | | |
| | (3) | menopause after | r 50 y | ears | (4) | won | nen over 40 ye | ars | |
| 78. | Gvn | ecomastia is a terr | m use | d for : | | | | | |
| , | (1) | male breast infe | | | (2) | fema | ale breast infec | tion | |
| | (3) | hypertrophy of | | breast | (4) | | ertrophy of fen | | st |
| 79. | ም ኤ - | shame of Campage | سے امن | famala malv | ia bui | ia . | | | |
| 79. | | shape of Gynaeco | | - | 15 0111 | | flat | (4) | hoort |
| | (1) | round | (2) | oval | | (3) | liat | (4) | heart |
| 80. | At to | erm pregnancy ut | erus e | enlarges fro | m 50 | gms to | o : | | |
| | (1) | 800 gms | (2) | 900 gms | | (3) | 850 gms | (4) | 950 gms |
| | | | | | | | | | |
| 81. | _ | ntening takes place | | | | | | | 24 |
| | (1) | 30 weeks | (2) | 32 weeks | | (3) | 34 weeks | (4) | 36 weeks |
| 82. | All | of the following a | re pos | sitive signs (| of pre | gnanc | y except : | | |
| | (1) | quickening | | | (2) | foeta | al heart sound | | |
| | (3) | movement of fo | etus | | (4) | Ame | enorrhea | | |
| 83. | App | proximate increase | in w | eight durin | g pres | gnanc | v is : | | |
| | (1) | 10.5 kg | (2) | 12.5 kg | 01 (| (3) | 11.5 kg | (4) | 13.5 kg |
| 84. | Brox | wnish colour loch | ia wh | ich occurs h | etwee | n 5 - | 9 davs is knov | n as: | |
| 01. | (1) | lochia serum | (2) | | | | lochia alba | (4) | lochiaserosa |
| | (^) | Total oca min | (~) | TOTTIM THE | _ • | (~) | Joseph dibu | (1) | |
| 85. | | ationship between wn as : | en the | long axis | of th | ne foe | tus and the l | ong axi | s of the uterus is |
| | (1) | presentation | (2) | attitude | | (3) | lie | (4) | position |

| 86. | | primigravida women the head nor | | |
|-----|------------|--|------------|--|
| | (1) (3) | 36 - 38 weeks of pregnancy 32 - 34 weeks of pregnancy | (2) (4) | * 0 - |
| | (3) | 52 - 54 Weeks of pregnancy | (4) | 50 - 52 weeks of pregnancy |
| 87. | | of the following are causes of abnormal | | - |
| | (1) | uterine dystocia | (2) | cervical dystocia |
| | (3) | foetal dystocia | (4) | pelvic dystocia |
| 88. | In vo | oluntary loss of products of concep | tion p | orior to 24 weeks of gestation is known as : |
| | (1) | spontaneous abortion | (2) | threatened abortion |
| | (3) | inevitable abortion | (4) | incomplete abortion |
| 89. | Hvp | eremesis Gravidarum starts before | : | |
| | (1) | 26 weeks of pregnancy | (2) | 24 weeks of pregnancy |
| | (3) | 22 weeks of pregnancy | (4) | 20 weeks of pregnancy |
| 90. | Who | m cold tamperature prohibits the | erozat- | and reproduction of bacteria is known as : |
| 90. | (1) | bacteriostatic (2) bactericida | - | (3) aerobic (4) anaerobic |
| | (1) | (2) vaccinos | 4, | (3) deroste (4) bracroste |
| 91. | | eonate weighing less than 2500 gms | at bir | th irrespective of the gestational age is referred |
| | as : | muo torma habri | (2) | love hinth regions habe |
| | (1) (3) | pre term baby very low birth weight baby | (2) (4) | low birth weight baby term baby |
| | (3) | very low butth weight baby | (4) | term vaby |
| 92. | | of the following are causes of hydr | oceph | alus except : |
| | (1) | obstruction in flow of fluid | | |
| | (2) | interference with absorption of fl | uid | |
| | (3) | leakage of CSF | | |
| | (4) | increased production of fluid | | |
| 93. | Und | escended testes is also known as : | | |
| | (1) | | (2) | cryptichidism |
| | (3) | cryptechidism | (4) | cryptuchidism |
| 94. | In ca | ase child is having phimosis with o | edem | a, it can be reduced by applying : |
| | (1) | cold compress | (2) | hot compress |
| | (3) | cold and hot compress | (4) | medicated hot compress |
| 95. | The | rapeutic diet advised for children v | with a | cute renal failure is : |
| - | (1) | low protein and high carbohydra | | |
| | (2) | low protein and high carbohydra | | d fat |
| | (3) | high protein and carbohydrate | | |
| | (4) | high protein and fat | | |

| 96. | All of | the following are at: | clinic | al feature of | a con | genita | al hypertrophic | pylorics | tenosis in chi | lare |
|------|------------|------------------------------------|---------|---------------|------------|-------------------|-------------------------------|-----------------|----------------|------|
| | (1) (3) | dehydration visible peristalsis | | | (2) (4) | | oilirus vomiting rgitation | 5 | | |
| 97. | Physi | iological jaundice | in te | rm babies su | ıbside | | | | | |
| | (1) | 5th day of birth | | | (2) | | ay of birth | | | |
| | (3) | 7 th day of birth | | | (4) | 8 th d | ay of birth | | | |
| 98. | Duri | ng phototherapy | the n | ursing meas | ures i | nclud | e all of the foll | owing e | xcept : | |
| | (1) | observation of h | - | | (2) | | sing up baby w | | ies | |
| | (3) | maintenance of | temp | erature | (4) | cove | ring eyes and g | genitalia | | |
| 99. | Pre te | erm babies are bo | rn be | fore : | | | | | | |
| | (1) | 37 weeks of gest | | | (2) | | eeks of gestati | | | |
| | (3) | 39 weeks of gest | ation | | (4) | 40 w | reeks of gestation | on | | |
| 100. | When | n an Infant turns | head | towards an | y obje | ect tha | at touches the o | cheek it | is known as | : |
| | (1) | rooting reflex | | | (2) | tonic | neck reflex | | | |
| | (3) | babinski's reflex | | | (4) | more | o reflex | | | |
| 101. | Apga | ar score of a newb | oorn | mmediately | after | the b | irth is checked | within : | | |
| | (1) | 1 minute | (2) | 2 minutes | | (3) | 3 minutes | (4) | 4 minutes | |
| 102. | Habi | tual ingestion of 1 | non-f | ood materia | l by a | maln | ourished child | is know | n as : | |
| | (1) | picro | (2) | piceous | | (3) | pico | (4) | pica | |
| 103. | All o | of the following as | re cor | ngenital abn | ormal | ities i | n new born <i>ex</i> | cept : | | |
| | (1) | Hydrocephalus | | | (2) | Men | ingitis | | | |
| | (3) | Tracheo - esoph | ageal | fistula | (4) | Spin | a - bifida | | | |
| 104. | All o | of the following as | re sig | ns and symp | ptom | of a c | hild with Mara | ismus <i>ex</i> | cept : | |
| | (1) | wasting of musc | les | | (2) | sunk | ken eyes | | | |
| | (3) | moon shaped fa | ce | | (4) | loos | e skin | | | |
| 105. | At b | irth normal head | circu | mference of | a nev | v bori | n is : | | | |
| | (1) | 28 to 30 cms | | | (2) | 30 to | o 32 cms | | | |
| | (3) | 33 to 35 cms | | | (4) | 36 to | o 38 cms | | | |
| 106. | Larg | ectile in high dose | es is 1 | nostly giver | n in or | ne of t | the following co | ondition | 5 | |
| | (1) | Schizophrenia | | | (2) | | genous depress | | | |
| | (3) | Endogenous der | oressi | on | (4) | Anx | riety | | | |
| | | | | | | | | | | |

| 107. | Nurs | se must observe one of the follo | wing sic | le effects in a patient who is on antipsychotics |
|------|-------|-----------------------------------|-----------|--|
| | (1) | Arrhythmia | (2) | Hypertension |
| | (3) | Extra Pyramidal symptom | (4) | Anorexia |
| 108. | Elect | tro Convulsive Therapy (ECT) is | most co | ommonly given in the following condition: |
| | (1) | Neurosis | (2) | Obsessive compulsive disorder |
| | (3) | Substance abuse | (4) | Depression |
| 109. | All c | of the following are types of Del | usions e | xcept : |
| | (1) | Grandiosity | (2) | Persecution |
| | (3) | Wrong perception | (4) | Ideas of reference |
| 110. | Whe | n an adolescent induces vomiti | ng after | taking adequate favourite food it is known as : |
| | (1) | Bulimia Nervosa | (2) | Anorexia Nervosa |
| | (3) | Obsessive Compulsive Neuros | is (4) | Neurosis |
| 111. | One | of the following is most commo | n symp | tom of opioid toxicity : |
| | (1) | constricted pupils | (2) | dialated pupils |
| | (3) | high fever | (4) | dysponea |
| 112. | Fear | of going outside to open space | is called | as: |
| | (1) | algo phobia | (2) | agro phobia |
| | (3) | acro phobia | (4) | amphi phobia |
| 113. | All | of the following are neurotic dis | orders e | xcept : |
| | (1) | Mania | (2) | Conversion disorders |
| | (3) | Exogenous depression | (4) | Phobia |
| 114. | Repe | etitive behaviour pattern perform | med in a | stereotype manner is known as : |
| | (1) | compulsion | (2) | obsession |
| | (3) | Repression | (4) | replication |
| 115. | ADI | HD stands for : | | |
| | (1) | Attention Deficit Hypoactive | Disorder | • |
| | (2) | Activity Deficit Hypoactive Di | sorder | |
| | (3) | Attention Deficit Hyperactive | Disorde | r |
| | (4) | Activity Deficit Hyperactive D | isorder | |
| 116. | All t | the following are therapeutic co | mmunic | ration techniques except: |
| | (1) | advising (2) observi | ng | (3) listening (4) silence |
| | | | | |

| 11/. | Prin | nary symptoms of sc | nizopnrenia ar | e an | except : | | |
|------|-------|-----------------------|------------------|--------|-------------------------|--------|-------------------|
| | (1) | Autistic thinking | | (2) | Ambivalence | | |
| | (3) | Auditory hallucina | tion | (4) | Appropriate respons | se . | |
| 118. | Patie | ent with mania are h | aving all of the | e folk | owing symptoms excep | pt: | |
| | (1) | elated and unstable | mood | (2) | sluggish and fatigue | | |
| | (3) | increased pressure | of speech | (4) | increased motor activ | vity | |
| 119. | Whe | en an individual expr | esses his failur | es an | d difficulties by blami | ng oth | ers is known as : |
| | (1) | repression (2 |) projection | | (3) sublimation | (4) | denial |
| 120. | Whe | en thoughts are too q | uick without a | ssoci | ation it is called : | | |
| | (1) | flight of ideas | | (2) | catalespy | | |
| | (3) | mannerisms | | (4) | clang association | | |
| | | | | | | | |

- o O o -