DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

T. B. C. : DS - 2015

Sefial No. 1785

Test Booklet Series



TEST BOOKLET

DENTAL SURGEON

Time Allowed : 3 Hours

(Maximum Marks : 200

: INSTRUCTIONS TO CANDIDATES :

- 1. IMMEDIATELY AFTER THE COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT THIS TEST BOOKLET **DOES NOT** HAVE ANY UNPRINTED OR TORN OR MISSING PAGES OR ITEMS ETC. IF SO, GET IT REPLACED BY A COMPLETE TEST BOOKLET OF THE SAME SERIES ISSUED TO YOU.
- 2. ENCODE CLEARLY THE TEST BOOKLET SERIES **A**, **B**, **C** OR **D**, AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK).
- 3. You have to enter your Roll No. on the Test Booklet in the Box provided alongside. DO NOT write anything else on the Test Booklet.
- 4. This Test Booklet contains 200 items (questions). Each item (question) comprises four responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question).
- 5. You have to mark (darken) all your responses (answers) ONLY on the separate Answer Sheet provided, by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet.
- All items (questions) carry equal marks. All items (questions) are compulsory. Your total
 marks will depend only on the number of correct responses (answers) marked by you in
 the Answer Sheet. There will be no negative marking for wrong answer.
- 7. Before you proceed to mark (darken) in the Answer Sheet the responses to various items (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per the instructions sent to you with your **Admission Certificate**.
- 8. After you have completed filling in all your responses (answers) on the Answer Sheet and after conclusion of the examination, you should hand over to the Invigilator the Answer Sheet issued to you. You are allowed to take with you the candidate's copy/second page of the Answer Sheet along with the Test Booklet after completion of the examination for your reference.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

1. Swan-neck deformity of the fingers Decreased serum ferritin (C) is seen in: Decreased Total Iron-binding (D) (A) Osteoarthritis Capacity (TIBC) Rheumatoid arthritis All are side effects of corticosteroids 6. (C) Osteomalacia except: (D) Osteoporosis (A) Myopathy Arthus type allergic reaction is a 2. Diarrhoea (B) feature of: Peptic ulcer disease (C) (A) Type I reactions (D) Osteoporosis (B) Type II reactions 7. Vitamin B12 deficiency may cause (C) Type III reactions all of the following except: Type IV reactions (D) (A) Optic neuritis Progression to chronic liver disease 3. Myelopathy (B) is seen most commonly with infection by: (C) Myopathy (A) Hepatitis A (D) Neuropathy (B) Hepatitis B Which of the following is the most 8. Hepatitis C (C) common site for a fracture Hepatitis D (D) associated with osteoporosis? Blood Coagulation Factor which is 4. (A) Femur not required in the intrinsic pathway (B) Hip for coagulation is: (C) Radius (A) Factor 9 (D) Vertebra (B) Factor 11 All are the causes of bilateral parotid 9. (C) Factor 7 enlargement except: (D) Factor 13 Diabetes mellitus (A) All the following are suggestive of 5. Systemic lupus erythromatosus (B) iron deficiency anemia except:

(A)

Koilonychia

Low reticulocyté response

(C)

(D)

HIV

Sarcoidosis

- 10. Which of the following is not a clinical feature of cerebellar disease?
 - (A) Nystagmus
 - (B) Hyperreflexia
 - (C) Hypotonia
 - (D) Tremors
- 11. All the following are Vitamin K-dependent coagulation factors except:
 - (A) Factor X
 - (B) Factor IX
 - (C) Factor III
 - (D) Factor VII
- 12. Excessive weight gain, menorrhagia, cold intolerance and weakness are suggestive of :
 - (A) Obesity
 - (B) Hypothyroidism
 - (C) Cushing Syndrome
 - (D) Anasarca
- 13. Screening test used for the diagnosis of Cushing syndrome is:
 - (A) ACTH stimulation test
 - (B) Plasma Cortisol
 - (C) Plasma FSH
 - (D) Plasma Aldosterone
- 14. Which serological marker is suggestive of highly infectious stage of acute Hepatitis B?
 - (A) HBsAg

- (B) Anti HBsAg
- (C) HBeAg
- (D) Anti HBeAg
- 15. Normal Serum Calcium, Phosphate and ALP are seen in :
 - (A) Osteomalacia
 - (B) Osteopetrosis
 - (C) Osteoporosis
 - (D) Osteonecrosis
- 16. Nasopharyngeal carcinoma mostly arises from the :
 - (A) Roof of nasopharynx
 - (B) Posterior wall
 - (C) Anterior wall
 - (D) Lateral wall
- 17. Highest incidence of nasopharyngeal carcinoma is seen in:
 - (A) Indian
 - (B) Chinese
 - (C) Americans
 - (D) Europeans
- 18. Which region is most commonly affected in actinomycosis?
 - (A) Thorax
 - (B) Liver
 - (C) Facio-cervical
 - (D) Liver

(3)

19.	A punched out edge is characteristic of which ulcer?		24		In tongue cancers, the least affected site is:		
	(A)	Tuberculosis		(A)	Ventral margin		
	(B)	Malignant		(B)	Dorsal margin		
	(C)	Rodent		(C)	Tīp		
	(D)	Syphilis		(D)	Lateral margin		
20.	Cys	tic hygroma is a :	25	. Am	eloblastoma most commonly		
	(A)	Dermoid cyst		dev	elops around :		
	(B)	Cavernous haemangioma		(A)	Incisors		
	(C)	Lymphangiectasia		(B)	Premolars		
	(D)	Sebaceous cyst		(C)	Canine		
21.	The	most common indication for the		(D)	Third molar		
	rem	oval of sublingual salivary gland	26	The	weakest part of mandible where		
	is:			frac	ture occurs is :		
	(A)	Ranula	•	(A)	Neck of condyle		
	(B)	Lymphangioma		(B)	Angle of mandible		
	(C)	Neoplasm		(C)	Midline		
.*	(D)	Stone		(D)	Canine fossa		
22.	Clef	t lip is ideally repaired at what	27.	Clef	t palate repair is ideally done at :		
	age	?		(A)	Birth		
	(A)	Soon after birth		(B)	1-6 months		
	(B)	6-9 weeks		(C)	7-18 months		
	(C)	10-24 weeks		(D)	24-36 months		
	(D)	25-36 weeks	28 .	The	most commonly involved bone in		
23.	Den	tigerous cyst develops around :	20.		ure of the face is:		
	(A)	Root		(A)	Nasal		
	(B)	Crown		(B)	Maxilla		
	(C)	Gum		(C)	Mandible		
:	(D)	Gingival Margin		(D)	Zygoma		
IB –	1A/34	ļ ((4)	` '	Contd.		

29.	Card	inoma of the lower alveolar		(C)	Globulins
	ridge	occurs around :		(D)	Prothrombin
	(A)	Incisor	34.	Exc	itation of baroreceptors in
	(B)	Canine		arte	ries reflexly causes :
•	(C)	Molar		(A)	Peripheral vasoconstriction
	(D)	Pre molar		(B)	Arterial pressure to decrease
30.	Sali	vary stones occurs most		(C)	Increase cardiac output
	com	monly in :		(D)	Vasovagal syncope
	(A)	Parotid	35.	Card	liac output is equal to the product
	(B)	Sublingual		of:	
	(C)	Submandibular		(A)	Stroke volume × venous return
	(D)	Submaxillary		(B)	Stroke volume × heart rate
31.	Bones, stones moans and groans			(C)	Venous return × heart rate
J1.		es to:		(D)	Stroke volume × cardiac index
		(A) Hyperparathyroidism		Pre-	optic area of hypothalamus is
	(B)			cond	cerned with :
	(C)	Hypoparathyroidism		(A)	Milk ejection
	(D)			(B)	Uterine contractility
	(0)	Hypothyroidism		(C)	Temperature regulation
32.		suture that maintains strength for		(D)	Memory
	long	est time is :	37.	REN	// sleep is associated with :
	(A)	Polyglycolic Acid (Dexon)		(A)	Normal heart rate
	(B)	Polyglactin (Vicryl)		(B)	Increased muscle tone
	(C)	Polydioxone (PDS)		(C)	Decreased respiratory rate
	(D)	Polyglyconate (Chromic catgut)		(D)	Extreme inhibition of peripheral
33.	Vitar	nin K is required by liver for the			muscles
	prod	uction of active :	38.	Sym	pathetic stimulation :
	(A)	Fibrinogen		(A)	Increases gastric motility
	(B)	Platelet		(B)	Causes Pupillary dilation
IB – 1A/34		(5)		(Turn over)	

(C) Decreases cardiac activity (B) Foetal haemoglobin (D) Vasodilation (C) Haemosiderin (D) Myoglobin 39. Which of the following best describes the Broca area of speech? 44. Deficiency of which one of the following does not lead to anemia? (A) Sensory speech area (A) Serum Iron (B) Controls emotions Serum Copper (B) (C) Located in temporal bone (C) Vitamin C (D) Motor speech area Serum Zinc (D) 40. Intentional tremor occurs in the 45. Most of the enzymes secreted in the disease of: human gastrointestinal tract are: (A) Cerebrum (A) Oxidases Basal ganglia (B) (B) Reductases (C) Cerebellum (C) Lyases (D) Spinal cord Hydrolases 41 Pain sensation travels through the: The primary function of the salivary 46. Posterior column (A) amylase is: Lateral spinothalamic tract (B) Hydrolysis of dietary starch (A) (C) Corticospinal tract Hydrolysis of bacterial cell wall (B) (D) Vestibulospinal tract (C) Hydrolysis of glycogen 42. ADH increases the permeability of (D) Hydrolysis of pectin distal tubule to: 47. Final common oxidative pathway (A) **Proteins** which integrates oxidative products (B) Amino acids of fats, proteins and carbohydrates (C) Glucose is: (D) Water (A) Gluconeogenesis (B) Glycolysis 43. Haemoprotein which binds to oxygen (C) Citric acid cycle most strongly is: Anaerobic glycolysis

(A) Adult haemoglobin

(D)

- 48. Largest amount of energy is produced by which of the following?
 (A) Oxidation of amino acids
 (B) Beta oxidation of saturated fatty acids
 (C) Beta oxidation of unsaturated fatty acid
 - (D) Anaerobic glycolysis
- 49. The substance that contributes maximally to the osmolality inside the cell is:
 - (A) Protein
 - (B) Phosphate
 - (C) Sodium
 - (D) Urea
- 50. The unique feature of Mitochondria is its:
 - (A) Actin
 - (B) Myosin
 - (C) Prothrombin
 - (D) DNA
- 51. Many substances are removed from the cell to outside by:
 - (A) Phagocytosis
 - (B) Pinocytosis
 - (C) Exocytosis
 - (D) Chemotaxis

- 52. Nerve fibre with slowest conduction is:
 - (A) A alpha
 - (B) A gamma
 - (C) Beta fiber
 - (D) C fibres
- 53. The superior meatus of the nose contains the opening of:
 - (A) Anterior ethmoidal air cells
 - (B) Middle ethmoidal air cells
 - (C) Posterior ethmoidal air cells
 - (D) Maxillary air sinus
- 54. The posterior auricular nerve is a branch of:
 - (A) Chorda tympani
 - (B) Facial
 - (C) Auriculo temporal
 - (D) Accessory nerve
- 55. The lateral wall of the infratemporal fossa is formed by:
 - (A) Lateral pterygoid plate
 - (B) Zygomatic arch
 - (C) Ramus of mandible
 - (D) Carotid sheath
- 56. The lymphatic from the tip of the tongue drains into:
 - (A) Submandibular lymph nodes
 - (B) Submental lymph nodes
 - (C) Jugulodigastric lymph nodes
 - (D) Jugulo omohyoid lymph nodes

57.	Which of the following is a branch of maxillary nerve? (A) Supratrochlear		62.	All are side effects of steroids except:		
			,	(A)	Myopathy	
	(A) (B)	Zygomaticotemporal		(B)	Diarrhoea	
	(C)	Infratrochlear		(C)	Peptic ulcer disease	
		External nasal		(D)	Osteoporosis	
	(D)		63.	Epin	ephrine causes all except :	
58.		taste sensation from the		(A)	Increase in blood glucose	
	-	erior one-third of the tongue is		` '	levels	
	(A)	ed by : Chorda tympani		(B)	Increase blood lactate	
7	(B)	Lingual nerve		(C)	Increases free fatty acids	
	(C)	Hypoglossal nerve		(D)	Increases triglycerides	
	(D)	Glossopharyngeal nerve	64.	Whi	ch of the following is an ester	
59.	Whic	ch of the following muscle assists		linke	ed local anesthetic?	
	in the elevation of mandible ?			(A)	Lidocaine	
	(A)	Medial pterygoid		(B)	Bupivacaine	
	(B)	Lateral pterygoid		(C)	Benzocaine	
	(C)	Digastric		(D)	Ropivacaine	
	(D)	Genohyoid	65.	Which of the following is heparin		
60.	The sensory root of the facial nerve			antagonist?		
	is kn	own as :		(A)	Warfarin	
	(A)	Lingual nerve		(B)	Vitamin K	
	(B)	Chorda tympani		` •		
	(C)	Nervous intermedius		(C)	Coumarin	
	(D)	Greater petrosal nerve		(D)	Protamine sulphate	
61.	Drug	causing hypocalcaemia is:	66.	Acut	e Asthma is treated with :	
	(A)	Lithium		(A)	Atropine	
	(B)	Phenytoin		(B)	Salbutamol	
	(C)	Thiazides		(C)	Digoxin	
	(D)	Vitamin A intoxication		(D)	Frusemide	

(8)

IB - 1A/34

Contd.

			•	
67.	Physical dependence is produced by:	72.	The most common site of intra-oral nevi is:	
	(A) Caffeine	•	(A) Hard palate	
	(B) Amphetamine	•	(B) Buccal mucosa	
	(C) Cocaine		(C) Gingiva	
	(D) Pethidine	:	(D) Lips	
68.	Intolerance is :			
	(A) Toxic effect of a drug in an individual at therapeutic doses	73.	The most common potentially malignant state of oral mucosa is:	
	(B) It indicates high threshold of		(A) Leukoplakia	
	the individual to the action of a		(B) Erythroplakia	
	drug		(C) Submucosal fibrosis	
	(C) Converse of the tolerance		(D) Keratoplakia	
	(D) All of the above	74.	The most common malignant	
69.	Endocrinal side effects of		neoplasm of the oral cavity is:	
	psychotropic drugs are:		(A) Squamous cell carcinoma	
	(A) Gynaecomastia		(B) Basal cell carcinoma	
	(B) Galactorrhea		(C) Malignant melanoma	
	(C) Both of the above		(D) Sarcoma	
	(D) None of the above			
70.	Drug of choice for febrile seizures:	75.	Unerrupted teeth is a common feature of:	
	(A) Diazepam		(A) Dentigerous cyst	
	(B) Phenytoin		(B) Odontogenic keratocyst	
	(C) Carbamate	a •	(C) Adenomatoid odontogenic	
	(D) Valproate		tumour	
71.	Gynotype in a patient with Turner		(D) All of the above	
c	syndrome is:			
	(A) 45XO	76.	Calcifying epithelial odontogenic	
	(B) 47XXY	•	tumour commonly occurs in:	
	(C) 46XX		(A) Mandibular Posterior area	

(9)

(B) Maxillary Posterior area

(Turn over)

(D) 47XYY

IB - 1A/34

- (C) Mandibular Anterior area
- (D) Maxillary Anterior area
- 77. The most common histological variant of Ameloblastoma is:
 - (A) Follicular
 - (B) Plexiform
 - (C) Acanthomatous
 - (D) Granular cell
- 78. The histological variant of ameloblastoma with highest recurrenence rate is:
 - (A) Follicular
 - (B) Acanthomatous
 - (C) Granular cell
 - (D) Basal cell
- 79. Smoking is associated with the genesis of which of the salivary gland tumor?
 - (A) Pleomorphic adenoma
 - (B) Warthin's tumor
 - (C) Necrotizing sialometaplasia
 - (D) Papillary cystadenoma
- 80. A tumor which arises in salivary gland tissue and entrapped in parotid lymph nodes during embryogenesis is:
 - (A) Pleomorphic adenoma
 - (B) Warthin's tumor
 - (C) Oncocytoma
 - (D) Mucinous adenocarcinoma

- 81. In which of the following condition the normal flora becomes pathogenic?
 - (A) Diabetes
 - (B) Leukemia
 - (C) AIDS
 - (D) All of the above
- 82. Following are the beneficial effects of the resident flora except:
 - (A) Synthesis of Vitamin K
 - (B) Production of Pencillinase
 - (C) Suppression of colonization by pathogen
 - (D) Bacteriocin production
- 83. Which of the following can be a congenital infection?
 - (A) AIDS
 - (B) Toxoplasmosis
 - (C) CMV
 - (D) All of the above
- 84. Following bacteria produces enterotoxin except:
 - (A) Staph aureus
 - (B) B. anthracis
 - (C) B. cereus
 - (D) V. cholerae
- 85. Which of the following disease is not acquired by ingestion?
 - (A) Salmonellosis
 - (B) Shigellosis
 - (C) Diptheria
 - (D) Cholerae

- 86. Following bacteria are invasive except:
 - (A) Shigella
 - (B) S. Typhi
 - (C) Vibiro Cholerae
 - (D) N. Meningitides
- 87. Which of the following is vector borne disease?
 - (A) Typhus
 - (B) Dengue
 - (C) Yellow fever
 - (D) All of the above
- 88. Plasmid mediated resistance was first detected in :
 - (A) Salmonella
 - (B) Shigella
 - (C) Proteus
 - (D) E. coli
- 89. Following are zoonotic disease except:
 - (A) Rabies
 - (B) Toxoplasmosis
 - (C) Hydatid
 - (D) Tetanus
- 90. Viral mediated transmission of genetic material in bacteria is known as:
 - (A) Transcription
 - (B) Translation
 - (C) Conjugation
 - (D) Transduction

- 91. Which is the cement with the highest solubility?
 - (A) $Ca(OH)_2$
 - (B) Silicate
 - (C) $Zn(PO_4)$
 - (D) GIC
- 92. Advantage of LED ligth is:
 - (A) Small and portable
 - (B) Low power consumption
 - (C) Minimal deterioration of light output
 - (D) All of the above
- 93. The angle between rake face and radial line is:
 - (A) Rake angle
 - (B) Clearance angle
 - (C) Blade angle
 - (D) Radial clearance
- 94. Eames technique provides for:
 - (A) Low mercury-alloy ratio
 - (B) High mercury-alloy ratio
 - (C) Easy waste management
 - (D) Good polishing
- 95. Which of the following is an estrogenic material?
 - (A) Bisphenol-A
 - (B) Poly acrylic acid
 - (C) Nickel
 - (D) DFG

96.	Simultaneous application of acid enamel and dentine is known as: (A) Total etch technique	to 101. Which of the following deciduous tooth does not resemble any of the tooth in permanent dentition?
	(B) Dry bonding	(A) A
	(C) Wet bonding	(B) B
	(D) Hybridization	(C) J
		(D) L
97.	Smear layer consists of :	102. Which of the following permanent
	(A) Enamel debris	tooth has a single antagonist in
	(B) Microorganism	occlusion?
,	(C) Dentin chips	(A) Maxillary third molar
	(D) All of the above	(B) Mandibular third molar
98.	Cement which is kindest to denta	(C) Mandibular canine
÷	pulp:	(D) Maxillary central incisor
	(A) GIC (B) Polycarboxylate	103. Which of the following tooth has mesial cusp slope greater than distal
	(C) Silicate	cusp slope ? (A) Maxillary first premolar
	(D) Resin cement	(B) Permanent mandibular canine
99.	Corrugated gold is a type of :	(C) Deciduous mandibular canine
	(A) Casting alloy	(D) Maxillary second premolar
	(B) DFG	104. Shape of occlusal aspect of maxillary
÷	(C) Technic alloy	permanent first molar is:
	(D) None of the above	(A) Trapezoidal
100		(B) Rectangle
100.	Elastic modulus of a materia indicates its:	(C) Square
	•	(D) Rhomboid
	(A) Rigidity	105. Which of the following is not a groove
	(B) Flexibility	pattern of mandibular second
	(C) Proportional limit	premolar?

(D) Viscosity

(À) Y

	(0)	11		(B)	Mandibular first premolar
	(C)	U		(C)	Permanent mandibular first
	(D)	M			molar
106	. Cel	ls concerned with resorption of		. (D)	Mandibular canine
		iduous teeth :	111	. Lec	ong's Premolar is :
	(A)	Osteoblast	, 1	(A)	Dens invaginatus
	(B)	Cementoclast		(B)	Dens evaginatus
	(C).	Odontoclast		(C)	Taurodontism
	(D)	Pericytes		(D)	Enamel hypoplasia
107	. Whi	ich of the following tooth contain	112	. Mos	st common benign tumor of oral
	sup	plementary grooves?		cav	ity is:
	(A)	Maxillary second premolar		(A)	Papilloma
	(B)	Mandibular second molar		(B)	Nevus
	(C)	Both (A) and (B)		(C)	Fibroma
	(D)	None of the above		(D)	Lipoma
108	. Coli	is seen in :	113.	. Ack	erman's tumor is :
	(A)	Buccal mucosa		(A)	Pleomorphic adenoma
	(B)	Inter dental papilla	,	(B)	Verrucous carcinoma
	(C)	Floor of mouth		(C)	Spindle cell carcinoma
	(D)	Hard palate		(D)	Adenolymphoma
109.		nelial rests of Malassez is seen	114.		ch of the following salivary gland
	in:				or is mimic mucocele?
•	(A)	Palate		(A)	Adenoid cystic carcinoma
	(B)	Gingiva		(B)	Salivary duct carcinoma
	(C)	Periodontal ligament		(C)	Mucoepidermoid carcinoma
	(D)	Dental pulp		(D)	Sebaceous carcinoma
110.	Mesi	olingual groove is seen in which	115.	Odo	ntoclastoma is :
	of the	e following teeth?		(A)	Giant cell tumor of bone
	(A)	Maxillary lateral incisor		(B)	Giant cell granuloma
IB-	1A/34	(13	3)		(Turn over)

Internal resorption of tooth 121. Drug of choice for the treatment of Actinomycosis is: (D) Hypercementosis Penicillin (A) 116. Mumps is caused by: Cephalosporin (B) Orthomyxovirus (A) Clotrimazole (C) (B) Adenovirus Metronidazole **Paramyxovirus** (D) Coxsackie virus 122. Moeller's Glossities is seen in : 117. Odontoclasia is: Iron deficiency Anaemia Linear enamel caries (A) Vitamin B6 deficiency **Tooth resorption** (B) (C) Vitamin B12 deficiency Odontogenic tumor (C) Thiamin deficiency Dry socket (D) 123. Life span of platelet is: 118. Bifid rib is seen in which of the (A) 6-8 days following syndrome? 12-14 days (A) Papillon Leferve syndrome (C) 60 days (B) Gorlin syndrome Sturge Weber syndrome 120 days (C) (D) (D) Ascher syndrome 124. Rushton bodies are present in: 119. Most common oral manifestation of (A) Periodontal cyst diabetes mellitus is: Dentigerous cyst Leukoplakia (A) (C) Odontogenic keratocyst Periodontal abscess Stafne's cyst (D) (C) ANUG 125. Muscle which depresses the (D) Hyperplastic candidiasis mandible is: 120. Alarm clock headache is: (A) Lateral Pterygoid Trigeminal neuralgia (A) Medial Pterygoid (B) (B) Temporal arteritis

Temporalis

Masseter

(C)

(D)

(C)

Sphenopalatine neuralgia

(D) Glossopharyngeal neuralgia

126.	Circu	moral pigmentation is seen in :		(B)	Henry Kempe
	(A)	Arsenic poisoning		(C)	Evangeline Jordan
	(B)	Lead poisoning		(D)	Addleston
	(C)	Peutz-Jaghers Syndrome		` '	
	(D)	Plummer-Vinson Syndrome	132.		erior bite plane is used for the
127.	Soap	bubble appearance is seen in :		corre	ection of :
	(A)	Osteosarcoma		(A)	Deep bite
	(B)	Osteoclastoma	•	(B)	Crowding
	(C)	Ewing's sarcoma		(C)	Anterior cross bite
	(D)	Multiple myeloma		(D)	Posterior cross bite
128.	_	used for the treatment of	133.	The	Daily dietary recommendation of
	serostomia is :			Calc	cium in children is :
	(A)	Atropine		(A)	600-800 mg
	(B)	Ephidrine Secondarine		(B)	800-1200 mg
	(C)	Scopolamine None of the above			-
	(D)			(C)	1200-1600 mg
129.	•	reflex is lost due to the paralysis	,	(D)	1600-2000 mg
	of : (A)	V th nerve	134	Nickel content in Stainless stee	
	(A) (B)	VII th nerve		crov	vn is :
		IX th nerve		(A)	17-19%
	(D)	XII th nerve		(B)	10-13%
130.	Dow	n's syndrome is trisomy of :	•	(C)	12-14%
	(A)	16 chromosome		(D)	11-12%
	(B)	18 chromosome	125	Ene	tein Pearls are found :
	(C)	21 chromosome	100	•	
	(D)	23 chromosome		(A)	On crest of ridges
131.	Pede	odontics treatment triangle was		(B)	Along mid palatine raphe
		n by:		(C)	Midline of mandible
	(A)	G. Z. Wright		(D)	On the buccal surface of incisor
IR -	1A/34	.	(15)		(Turn over)

136.	Perio	od of mixed dentition growth		(C)	Maxilla-5mm, Mand-7mm
	spurt	in boys:		(D)	Maxilla-5mm, Mand-3mm
	(A)	10-11 years	141.	EH /	Angle introduced his classi-
	(B)	7-10 years			on of malocclusion in :
,	(C)	12-13 years		(A)	1988
	(D)	8-11 years		(B)	1899
137.	Whic	ch of the following GIC is using		(C)	1901
•	as Pi	t and Fissure Sealent?		(D)	1801
	(A)	Type 6	140		most important augsbandrasis for
	(B)	Type 8	142.		most important synchondrosis for growth modification is:
٠.	(C)	Type 7		(A)	Spheno ethmoidal
	(D)	Type 9		(B)	Spheno occipital
138.	The	first evidence of calcification of		(C)	Inter sphenoidal
	Primary teeth begins approximately			(D)	Inter maxillary
	at:				
	(A)	11 weeks in utero	143.	Mon	nent of a force is:
	(B)	14 months in utero		(A)	Force × perpendicular from
	(C)	16 months in utero	,		centre of rotation
	(D)	6 months in utero		(B)	Force × perpendicular from
139.	Clas	ssical Conditioning was			centre of resistance
	prop	osed by:	ż	(C)	Area under the stress strain
	(A)	Skinner			graph
	(B)	Pavlov	•	(D)	Force × range
	(C)	Peaget	144.	Third	d order bends are :
	(D)	Hassel		(A)	In-out bends
140.	Incis	al Liability is :		(B)	Tip back bends
	(A)	Maxilla-3mm, Mand-5mm		(C)	Torquing bends
	(B)	Maxilla-7mm, Mand-5mm		(D)	Anti-rotation bends
IB-	1A/34	(16	i)		Contd.

145. A	Anterior bolton's ratio is :	150.	. Rev	verse orthodontics refers to :
(/	A) 77.2%		(A)	Repeated orthodontic treat-
(1	B) 91.3%			ment
	C) 88.2%		(B)	Camouflage treatment
-	D) 91.7%		(C)	De-compensation
	/ /luscle affected in torticollis :		(D)	Treatment using lingual
		•		appliances
	A) Lateral pterygoid	151.	June	ctional epithelium shows rete peg
	B) Sternocleidomastoid		form	nation in :
• •	C) Buccinators		(A)	Stage I gingivitis
(L	D) Temporalis		(B)	Stage II gingivitis
147. N	lidline shift with unilateral posterio	r	(C)	Stage III gingivitis
	ross bite may be due to a change	Э	(D)	Stage IV gingivitis
in		152.	Reu	nion of connective with root
	A) Pitch			ace a root surface that has been
(E			path	ologically exposed :
	رک) Yaw		(A)	Reattachment
(E	D) None of the above		(B)	New attachment
148. Te	enth theorem of retention was pu	t	(C)	Regeneration
, / fo	orth by:		(D)	Repair
(A	A) R. Riedel	153.	If ha	If of the crown along with root is
(E	3) W. R. Proffit			arated or one half is removed, it
(0	C) A. Lundstrom		is ca	lled:
(0	D) R. E. Moyers		(A)	Radisection
149. N	ormal naso labial angle necessi	_	(B)	Hemisection
ta	tes the extraction of :		(C)	Apexification
(A	N) Upper first premolar		(D)	Coronosection
(E	B) Upper second premolar	154.	Most	abundant immunoglobulin in the
(C	C) Upper incisor		seru	m is :
(C) None of the above		(A)	lgG
IB – 1A	/34	(17)		(Turn over)

	(B)	lgA			(C)	4 months
. ,	(C)	lgM			(D)	2 months
	(D)	IgE		160.	The	e microorganism found in higher
155	. Whi	ch of the following grafts has th				s in pregnancy gingivitis :
		prognosis?			(A)	P. gingivalis
-	(A)	Xenograft			(B)	P intermedia
	(B)	Autograft			(C)	Capnocytophaga
	(C)	Allograft			(D)	A. viscosus
	(D)	Homograft		101		
156.	. Ram	ping can be done for a :	,	161.		h ml of 2% lignocaine contains :
	(A)	One wall defect			(A)	0.2 mg of lignocaine
	(B)	Two wall defect			(B)	2 mg of lignocaine
	(C)	Three wall defect			(C)	10 mg of lignocaine
	(D)	Zero wall defect			(D)	20 mg of lignocaine
157	Halin	neters are used in :	1	62.	Whi	ch of the following mandibular
107.	(A)	Oral malodour			teet	h may be removed without pain
	(A) (B)	Tooth mobility				rthe administration of IANB and
					lingu	ual block ?
	(C)	Pathologic migration	. •	• .	(A)	All molars
	(D)	Pulp testing			(B)	Both premolars and molars
158.		cept of full mouth disinfection	1		(C)	All teeth in lower jaw
		out forward by :			(D)	First premolar and canine
	` ,	Miller	1	63.	Follo	owing structures are present in
	` '	Glickman			the s	ubmandibular space except :
		William			(A)	Deep part of submandibular
		Quirynen				gland
159.		nelisation of gingivectomy	/		(B)	Facial artery
111		d is completed by:			(C)	Mylohyoid nerve and vessels
		4 days			(D)	Superficial part of sub-
	(B)	1 month				mandibular gland
IB-	A/34		(18)	_		Contd.

- 164. Surgical emphysema may develop immediately after:
 - (A) Cutting the tooth with airotor
 - (B) Cutting the tooth with micromotor
 - (C) Cutting the bone with micromotor
 - (D) Cutting the bone with airotor
- 165. Metronidazole is a:
 - (A) Quinolone
 - (B) Anti-fungal
 - (C) Imidazole derivative
 - (D) Antiviral
- 166. Which of the following artery does not accompany the nerve throughout its course?
 - (A) Lingual
 - (B) Infra orbital
 - (C) Inferior alveolar
 - (D) Posterior superior alveolar
- 167. Secondary bony callus formation takes approximately:
 - (A) 1-2 weeks
 - (B) 3-4 weeks
 - (C) 4-6 weeks
 - (D) 6-8 weeks
- 168. Which of the following is DANGER SPACE?
 - (A) Pterygo mandibular
 - (B) Lateral pharyngeal

- (C) Superficial and deep temporal
- (D) Space between alar and prevertebral fascia
- 169. Weber Fergussion incision is used for:
 - (A) Mandibulectomy
 - (B) Maxillectomy
 - (C) Condylectomy
 - (D) Rhinoplasty
- 170. Cranial nerve passing through the centre of the cavernous sinus is:
 - (A) Optic nerve
 - (B) Ophthalmic nerve
 - (C) Occulo motor nerve
 - (D) Abducent nerve
- 171. Articular disc of TMJ is:
 - (A) Fibro cartilage
 - (B) Hyaline cartilage
 - (C) Elastic tissue
 - (D) TMJ ligament
- 172. Four clotting factors synthesized in liver are:
 - (A) III, V, VII and VIII
 - (B) II, VII, IX and X
 - (C) II, VI, IX and XII
 - (D) II, V, VII and IX
- 173. Which of the following is efficacious modality in diagnosis of salivary gland neoplasm?
 - (A) Incisional biopsy

Excision biopsy (B) Lower member (B) (C) Fine needle aspiration cytology (C) No condyle (D) MRI scan (D) Fixed condyle 174. 1: 1000 adrenaline ampule contains: 179. The quality that distinguishes one (A) 0.1 mg/ml from the other is: (B) 1 mg/ml (A) Hue 10 mg/ml (C) Chroma 0.01 mg/ml Value 175. The target of gow gates technique Saturation (D) is: 180. In laminate technique of impression (A) Coronoid notch making: (B) Sigmoid notch (A) Alginate and agar are used Mandibular foramen (B) Only alginate is used (D) Neck of condyle Elastomers are used 176. "Enter bolus-exit balance" was stated Only agar by: (D) (A) Harold Ortman 181. Central incisors are generally (B) Sheppard anterior to the line bisecting incisive (C) Stewart pappilae by: (D) Hudson (A) 8-10 mm 177. The estimated width of maxillary (B) 14-15 mm central incisor is: (C) 2-6 mm 1/16th bizygomatic width (A) 11-20 mm (D) 1/16th interpupillary distance (B) 182. Advantages of an over denture: 1/8th bizygomatic width (C) Preservation of ridge (A) (D) None of the above Support (B) 178. In arcon articulators the condylar Retention element is located in: All of the above (D) (A) Upper member

(20)

IB - 1A/34

Contd.

183	. Fac	ebow transfer records:		(C)	Inhibitor
	(A)	Orientation relation		(D)	Plasticizer
	(B)	Special relation	188	. Ala	inate is made dust free by the
	(C)	Vertical relation			lition of:
	(D)	Tentative centric relation		(A)	Glycerine
184	. Idei	ntify the correct hanau formula	3	(B)	Calcium sulphate
		ch gives the lateral condyla		(C)	Sodium phosphate
	guio	lance angle :		(D)	Calcium chloride
	(A)	L = H/8 + 12	189	. Res	st seat can be prepared :
	(B)	L = 8/H + 12		(A)	Only on tooth surface
	(C)	L = H/8 * 12		(B)	In amalgam restoration
	(D)	L = H/8 – 12		(C)	In cast gold restoration
185.	. Inst	rument used to locate ptery	-	(D)	All of the above
	gom	axillary notch :	190.	Hea	I raising movement of a RPD is
	(A)	Boby gauge			vented by:
	(B)	Divider	:	(A)	Occlusal rest
	(C)	Screw gauge	V	(B)	Minor connector
	(D)	T-burnisher	1 4 t 1	(C)	Direct retainer
186.	The	distal border of mandibular	,	(D)	Indirect retainer
	dent	ure should cover :	191.	India	cator in Snyder Test is:
-	(A)	Ant. $\frac{1}{3}$ rd of retromolar pad		(A)	Diazoresorcinol
	(B)	Ant. $\frac{2}{3}$ rd of retromolar pad		(B)	Methyl red
	(C)	Should not cover the retromolar		(C)	Glucose
		pad		(D)	Bromocresol green
	(D)	Hamullar notch	192	Firet	artificial water fluoridation of
187.	In ac	rylic resin, hydroquinone act as :	102.		er supply was done in :
	(A)	Activator		(A)	London
	(B)	Initiator	ty to give	(B)	New York
IB –	1A/34	, 	(21)	-	(Turn over)

- (C) Grand Rapids
- (D) Ontario, Canada
- 193. Specificity of a test denotes:
 - (A) True Positive
 - (B) True Negative
 - (C) False Positive
 - (D) False Negative
- 194. The coverage of the geographic area through dental clinics can be depicted by the following way:
 - (A) Frequency polygon
 - (B) Cartogram
 - (C) Histogram
 - (D) Pie diagram
- 195. An experiment in which the experimental subjects cannot distinguish the experimental conditions is called as:
 - (A) Two Tailed Testing
 - (B) Single Blind experiment
 - (C) Double Blind experiment
 - (D) Retrospective experiment
- 196. Pit and fissure sealant is a form of:
 - (A) Primary prevention
 - (B) Secondary prevention

- (C) Tertiary prevention
- (D) Primordial prevention
- 197. Shoe leather survey was carried out by:
 - (A) Churchill H. V.
 - (B) Frederick McKay
 - (C) Trendley H. Dean
 - (D) Greene Vardiman Black
- 198. Example of a composite index is:
 - (A) CPITN
 - (B) DMF
 - (C) TNM
 - (D) Periodontal
- 199. Occurrence of disease clearly in excess of expected occurrence:
 - (A) Endemic
 - (B) Pandemic
 - (C) Sporadic
 - (D) Epidemic
- 200. Which of the following is a measure of dispersion?
 - (A) Mean
 - (B) Median
 - (C) Mode
 - (D) Standard deviation

SPACE FOR ROUGH WORK